LebanonTurf

Country Club®

Quick aerification recovery Speeds up root establishment Increases turf density

GRANULAR FERTILIZER + mycorrhizal fungi

GUARANTEED ANALYSIS	
Total Nitrogen (N)	3.0%
1.50% Ammoniacal Nitrogen	
1.50% Water Insoluble Nitrogen*	
Available Phosphate (P ₂ 0 ₅)	3.0%
Soluble Potash (K ₂ 0)	4.0%
Calcium (Ca)	4.0%
Magnesium (Mg)	1.5%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.01% Water Soluble Iron (Fe)	

Derived From: Wheat bran, poultry compost, feather meal, potassium sulfate, diammonium sulfate, ammonium sulfate, kelp meal, ferrous sulfate and ferric oxide. *1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

0.05% Amino Acid (Glyd
0.02% Myo-Inositol
0.01% Thiamine (Vitam
***** *****

ALSO CONTAINS NON PLANT FOOD INGREDIENTS:	Mycorrhizal fungi (#viable pr	opagules/gram)
0.50% Humic Acids	Rhizophagus irregularis	1.2
0.15% Ascorbic Acid (Vitamin C)	Rhizophagus clarus	1.2
0.05% Amino Acid (Glycine)	Rhizophagus aggregatus	0.6
0.02% Myo-Inositol	Rhizophagus diaphanus	1.2
0.01% Thiamine (Vitamin B₁)	Funneliformis mosseae	1.2
	Funneliformis monosporus	0.6
	Septoglomus deserticola	1.2
****	Claroideoglomus etunicatum	1.2
	Claroideoglomus claroideum	1.2

GUARANTEED MINIMUM ANALYSIS


Total Nitrogen (N)	3.0%
1.50% Ammoniacal Nitrogen	
1.50% Water Insoluble Nitrogen*	
Available Phosphate (P205)	3.0%
Soluble Potash (K ₂ 0)	4.0%
Calcium (Ca)	4.0%
Magnesium (Mg)	1.5%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.01% Water Soluble Iron (Fe)	

Derived From: Wheat bran, poultry compost, feather meal, potassium sulfate, diammonium sulfate, ammonium sulfate, kelp meal, ferrous sulfate and ferric oxide. *1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS: Mycorrhizal fungi (#viable propagules/gram)

Rhizophagus irregularis

0.15% Ascorbic Acid (Vitamin C)	Rhizophagus cla
0.05% Amino Acid (Glycine)	Rhizophagus ag
0.02% Myo-Inositol	Rhizophagus dia
0.01% Thiamine (Vitamin B ₁)	Funneliformis m
	Funneliformis m
	Septoglomus de
	Claroideoglomus
	Claroideoglomus



NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

0.50% Humic Acids

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

Lesco (letter/Numeric)	L/24	M/26
PennMulch HVO	P	Q
ProScape SS	P	Q
Spyker	6	6.5
Vicon (all models)	34	40
	Spyker Vicon (all models)	Spyker 6

_ebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation 1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

GUARANTEED ANALYSIS

Total Nitrogen (N) 3.0%

3.0%
4.0%
4.0%
1.5%
3.0%
1.0%

Derived From: Wheat bran, poultry compost, feather meal, potassium sulfate, diammonium sulfate, ammonium sulfate, kelp meal, ferrous sulfate and ferric oxide. *1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS:

0.50% Humic Acids	
0.15% Ascorbic Acid (Vitamin C)	
0.05% Amino Acid (Glycine)	
0.02% Myo-Inositol	
0.01% Thiamine (Vitamin B₁)	



wycorrnizai	tungi	(#viable	propaguies/grai	n)
Dhizophoguo	irrogul	orio	10	

Rhizophagus irregularis	1.2
Rhizophagus clarus	1.2
Rhizophagus aggregatus	0.6
Rhizophagus diaphanus	1.2
Funneliformis mosseae	1.2
Funneliformis monosporus	0.6
Septoglomus deserticola	1.2
Claroideoglomus etunicatum	1.2
Claroideoglomus claroideum	1.2

F699

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	3.0%
1.50% Ammoniacal Nitrogen	
1.50% Water Insoluble Nitrogen*	
Available Phosphate (P ₂ 0 ₅)	3.0%
Soluble Potash (K ₂ 0)	4.0%
Calcium (Ca)	4.0%
Magnesium (Mg)	1.5%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.01% Water Soluble Iron (Fe)	

Derived From: Wheat bran, poultry compost, feather meal, potassium sulfate, diammonium sulfate, ammonium sulfate, kelp meal, ferrous sulfate and ferric oxide. *1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS: Mycorrhizal fungi (#viable propagules/gram)

0.50% Humic Acids	Rhizophagus irregularis	1.2
0.15% Ascorbic Acid (Vitamin C)	Rhizophagus clarus	1.2
0.05% Amino Acid (Glycine)	Rhizophagus aggregatus	0.6
0.02% Myo-Inositol	Rhizophagus diaphanus	1.2
0.01% Thiamine (Vitamin B ₁)	Funneliformis mosseae	1.2
	Funneliformis monosporus	0.6
	Septoglomus deserticola	1.2
	Claroideoglomus etunicatum	1.2
	Claroideoglomus claroideum	12

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

Spreader	8#/1,000 sq. ft.or 3.9 kg/100m² (covers up to 5,000 ft² of lawn) (covers up to 568m² of lawn)	10#/1,000 sq. ft. or 4.8 kg/100m² (covers up to 5,000 ft² of lawn) (covers up to 454m² of lawn)	Spreader	8#/1,000 sq. ft.or 3.9 kg/100m² (covers up to 5,000 ft² of lawn) (covers up to 568m² of lawn)	10#/1,000 sq. ft. or 4.8 kg/100m² (cover up to 5,000 ft² of lawn) (covers up to 454m² of lawn)
LebanonTurf	6	6.5	Lesco (letter/Numeric)	L/24	M/26
Andersons AccuPro	M	0	PennMulch HVO	Р	Q
Earthway Rotary	20	24	ProScape SS	Р	Q
Gandy	32	36	Spyker	6	6.5
Lely	6.5 II	7	Vicon (all models)	34	40
These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.					



TURF APPLICATIONS

Apply 8 lbs. per 1,000 sq. ft. (4kg per 100m²) **Sod / Seed:** Apply 10 lbs. per 1,000 sq. ft. (5kg per 100m²)

In Florida - Apply 8 lbs. per 1,000 sq. ft. Maximum of 2 applications per year)

Divot Mix: Add 25 lbs. per cubic yard of divot mix

PLANTINGS

Mix thoroughly with the soil being used to fill the hole where being planted.

1/2 lb. (1 cup or 227g) for every 1" diameter of trunk

Shrubs: One gallon (3.7 liters) containers – ½ lb. (1/2 cup or 110g)

Five gallon (18.9 liters) containers – 1 lb. (2 1/4 cups or 450g)

When used as directed. Country Club Root Reviver will supplement your regular fertilizer program by providing up to .30 lbs. of nitrogen per 1,000 sq. ft. (In Florida - .24 lbs. of nitrogen per 1,000 sq. ft.)

WARNING



Hazard Statements

Causes skin irritation.

May cause respiratory irritation.

Causes serious eye irritation.

Causes eye irritation

Precautionary Statements

Wash hands and exposed skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection.

If in eyes, rinse with water and seek medical attention if symptoms persist.

Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Store locked up.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container properly as required by law for waste pesticides.

Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;
 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended.
- 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
- or 3) Re-establishing or repairing a turf area.
- *In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Not for use on Food Crops.