

LebanonTurf

# Country Club<sup>®</sup>

## Blended Fertilizer

# 8-4-24



With **HUMIC MAX** & **MESA**

### GUARANTEED ANALYSIS

Total Nitrogen (N)	8%
1.8% Ammoniacal Nitrogen	
1.3% Water Insoluble Nitrogen*	
2.6% Urea Nitrogen	
2.3% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	24%
Sulfur (S)	9.0%
9.0% Combined Sulfur (S)	

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.  
\*3.6% Slowly Available Nitrogen from Methylene Ureas.

U.S.A

NON-PLANT FOOD INGREDIENTS  
Humic acids (derived from Leonardite)

### GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	8%
1.8% Ammoniacal Nitrogen	
1.3% Water Insoluble Nitrogen*	
2.6% Urea Nitrogen	
2.3% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	24%
Sulfur (S)	9.0%
9.0% Combined Sulfur (S)	

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.  
\*3.6% Slowly Available Nitrogen from Methylene Ureas.

CANADA

NON-PLANT FOOD INGREDIENTS  
Humic acids (derived from Leonardite)

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Lbs N/1,000 Sq. Ft.	Lbs. P/1,000 Sq. Ft.	Lbs. K/1,000 Sq. Ft.	Lbs Humic Acids/1,000 Sq. Ft.	Lbs Fert/1,000 Sq. Ft.	Lbs Required for One Acre	Sq. Ft. Coverage per bag
0.9	0.25	1.5	0.5	6.2	270	8,050
0.5	0.13	0.75	0.25	3.1	135	16,100

**Application Rates:**  
To feed at the rate of 1.5 lb. Potash (K<sub>2</sub>O) per 1,000 sq. ft., apply this product at 6.2 lbs per 1,000 sq. ft. or 270 lbs. per acre. To feed at the rate of 1 lb. Potash (K<sub>2</sub>O) per 1,000 sq. ft., apply this product at 3.1 lbs per 1,000 sq. ft. or 135 lbs. per acre.

Spreader	1.5 # K <sub>2</sub> O /1,000 sq. ft. ½ # K <sub>2</sub> O /1,000 sq. ft.	4%	4%
Lebanon Turf		4%	4%
Andersons AccuPro	M	i	i
Earthway Rotary	16	15	15
Gandy	28	25	25
Lely	6½ II	5 II	5 II
Lesco (letter/numeric)	J/21	i/16	i/16
PennMulch HVO	M	K	K
ProScape SS	M	K	K
Spyker	4%	4%	4%
Vicon	26	23	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

**In Maryland, New Jersey, New York and Virginia:** This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory\* performed no more than 3 years before the application;
- 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
- or 3) Re-establishing or repairing a turf area.

\*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

**In New Hampshire:** This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

### Hazard Statements

Causes skin irritation.  
May be harmful if inhaled.  
May cause respiratory irritation.  
Causes eye irritation

### Warning



### Precautionary Statements

Wash hands and exposed skin thoroughly after handling.  
Wear protective gloves/protective clothing/eye protection/face protection.  
If in eyes, rinse with water and seek medical attention if symptoms persist.  
Take off contaminated clothing and wash before reuse.  
IF ON SKIN: Wash with plenty of soap and water.  
IF INHALED: Call > POISON CENTER or doctor/physician if you feel unwell.  
If skin irritation occurs: Get medical advice/attention.  
If eye irritation persists: Get medical advice/attention.  
Store locked up.  
Store in a well-ventilated place.  
Keep container tightly closed.  
Dispose of contents/container properly as required by law for waste pesticides.  
Keep out of reach of children  
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground

### WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Country Club, PennMulch, ProScape and the Lebanon Diamond are registered trademarks of Lebanon Seaboard Corporation.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

**Application Rates:**  
To feed at the rate of 0.48 kg Potash (K<sub>2</sub>O) Equivalent to 3.2 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Potash (K<sub>2</sub>O) Equivalent to 2.1 kg of product per 100 sq. meters.

Spreader	0.48 kg K <sub>2</sub> O/100 m <sup>2</sup> .	0.24 kg K <sub>2</sub> O/100 m <sup>2</sup> .
Lebanon Turf	4%	4%
Andersons AccuPro	M	i
Earthway Rotary	16	15
Gandy	28	25
Lely	6½ II	5 II
Lesco (letter/numeric)	J/21	i/16
PennMulch HVO	M	K
ProScape SS	M	K
Spyker	4%	4%
Vicon	26	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699 22-37132 i/20

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

LebanonTurf

For technical assistance or more information about our products visit [www.LebanonTurf.com](http://www.LebanonTurf.com)


**Lebanon Seaboard Corporation**  
1600 E. Cumberland St • Lebanon, PA 17042  
800-233-0628 • (717) 273-1685

**50 Lbs. (22.7 kg) Net Weight**

# 8-4-24



With



## GUARANTEED ANALYSIS

Total Nitrogen (N) .....	8%
1.8% Ammoniacal Nitrogen	
1.3% Water Insoluble Nitrogen*	
2.6% Urea Nitrogen	
2.3% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	24%
Sulfur (S) .....	9.0%
9.0% Combined Sulfur (S)	

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.  
\*3.6% Slowly Available Nitrogen from Methylene Ureas.

**U.S.A**

### NON-PLANT FOOD INGREDIENTS

Humic acids (derived from Leonardite)

## GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N) .....	8%
1.8% Ammoniacal Nitrogen	
1.3% Water Insoluble Nitrogen*	
2.6% Urea Nitrogen	
2.3% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	24%
Sulfur (S) .....	9.0%
9.0% Combined Sulfur (S)	

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.

\*3.6% Slowly Available Nitrogen from Methylene Ureas.

**CANADA**

### NON-PLANT FOOD INGREDIENTS

Humic acids (derived from Leonardite)

**Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.**

Lbs N/1,000 Sq. Ft.	Lbs. P/1,000 Sq. Ft.	Lbs. K/1,000 Sq. Ft.	Lbs Humic Acids/1,000 Sq. Ft.	Lbs Fert/1,000 Sq. Ft.	Lbs Required for One Acre	Sq. Ft. Coverage per bag
0.9	0.25	1.5	0.5	6.2	270	8,050
0.5	0.13	0.75	0.25	3.1	135	16,100

### Application Rates:

To feed at the rate of 1.5 lb. Potash (K<sub>2</sub>O) per 1,000 sq. ft., apply this product at 6.2 lbs per 1,000 sq. ft. or 270 lbs. per acre. To feed at the rate of 1 lb. Potash (K<sub>2</sub>O) per 1,000 sq. ft., apply this product at 3.1 lbs per 1,000 sq. ft. or 135 lbs. per acre.

### SUGGESTED SPREADER SETTINGS

Spreader	1.5 # K <sub>2</sub> O /1,000 sq. ft. ½ # K <sub>2</sub> O /1,000 sq. ft.	4¼	4¼
Lebanon Turf	4¼	4¼	
Andersons AccuPro	M	i	
Earthway Rotary	16	15	
Gandy	28	25	
Lely	6½ II	5 II	
Lesco (letter/numeric)	J/21	i/16	
PennMulch HVO	M	K	
ProScape SS	M	K	
Spyker	4¼	4¼	
Vicon	26	23	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

**In Maryland, New Jersey, New York and Virginia:** This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory\* performed no more than 3 years before the application;
- 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
- or 3) Re-establishing or repairing a turf area.

\*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

**In New Hampshire:** This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

### Hazard Statements

Causes skin irritation.  
May be harmful if inhaled.  
May cause respiratory irritation.  
Causes eye irritation

## Warning



### Precautionary Statements

Wash hands and exposed skin thoroughly after handling.  
Wear protective gloves/protective clothing/eye protection/face protection.  
If in eyes, rinse with water and seek medical attention if symptoms persist.  
Take off contaminated clothing and wash before reuse.  
IF ON SKIN: Wash with plenty of soap and water.  
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.  
If skin irritation occurs: Get medical advice/ attention.  
If eye irritation persists: Get medical advice/attention.  
Store locked up.  
Store in a well-ventilated place.  
Keep container tightly closed.  
Dispose of contents/container properly as required by law for waste pesticides.  
Keep out of reach of children  
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground

### WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Country Club, PennMulch, ProScape and the Lebanon Diamond are registered trademarks of Lebanon Seaboard. Corporation.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

### Application Rates:


To feed at the rate of 0.48 kg Potash (K<sub>2</sub>O)  
Equivalent to 3.2 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Potash (K<sub>2</sub>O)  
Equivalent to 2.1 kg of product per 100 sq. meters.

### SUGGESTED SPREADER SETTINGS

Spreader	0.48 kg K <sub>2</sub> O/100 m <sup>2</sup> .	0.24 kg K <sub>2</sub> O /100 m <sup>2</sup> .
Lebanon Turf	4¼	4¼
Andersons AccuPro	M	i
Earthway Rotary	16	15
Gandy	28	25
Lely	6½ II	5 II
Lesco (letter/numeric)	J/21	i/16
PennMulch HVO	M	K
ProScape SS	M	K
Spyker	4¼	4¼
Vicon	26	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699 **22-37132** i/20

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>