

LebanonTurf



Country Club[®]

Blended Fertilizer

25-0-15

MESA
PATENTED

GUARANTEED ANALYSIS

Total Nitrogen	25%
4.3% Ammoniacal Nitrogen	
2.7% Water Insoluble Nitrogen*	
13.4% Urea Nitrogen	
4.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	15%
Sulfur (S)	5.0%
5.0% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.1% Water Soluble Iron	
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
*7.3% Slowly Available Nitrogen from Methylene Ureas.	
Chlorine (Cl)	11.0%

U.S.A.

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen	25%
4.3% Ammoniacal Nitrogen	
2.7% Water Insoluble Nitrogen*	
13.4% Urea Nitrogen	
4.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	15%
Sulfur (S)	5.0%
5.0% Combined Sulfur (S)	
Iron (Fe) Actual	1.0%
0.1% Water Soluble Iron	
Chlorine (Cl)	11.0%
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
*7.3% Slowly Available Nitrogen from Methylene Ureas.	

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

WARNING

Hazard Statements:
Causes eye irritation.
May be harmful if inhaled

Precautionary Statements for handling:
Wash hands and exposed skin thoroughly after handling.
If in eyes, rinse with water and seek medical attention if symptoms persist.
IF INHALED: Call - POISON CENTER or doctor/physician if you feel unwell.
If eye irritation persists: Get medical advice/attention.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen turf.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.6 lbs per 1,000 sq. ft. or 156.5 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.0 lbs per 1,000 sq. ft. or 87 lbs. per acre.

Spreader	SUGGESTED SPREADER SETTINGS	
	0.9#N/1,000 sq. ft.	0.5#N/1,000 sq. ft.
Lebanon Turf	4½	4
Andersons AccuPro	L	H
Earthway Rotary	15	14
Gandy	27	24
Lely	5½ II	4 II
Lesco (letter/numeric)	i/19	G/15
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	4½	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

Application Rates:

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.
To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 0.98 kg of product per 100 sq. meters.

Spreader	SUGGESTED SPREADER SETTINGS	
	0.43 kg N/100 m ² .	0.22 kg N/100 m ² .
Lebanon Turf	4½	4
Andersons AccuPro	L	H
Earthway Rotary	15	14
Gandy	27	24
Lely	5½ II	4 II
Lesco (letter/numeric)	i/19	G/15
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	4½	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-37090

i/19

MESA[®], PennMulch[®] and ProScape[®] are registered trademarks of Lebanon Seaboard Corporation.

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
1600 E. Cumberland St • Lebanon, PA 17042
800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight

25-0-15

MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen	25%
4.3% Ammoniacal Nitrogen	
2.7% Water Insoluble Nitrogen*	
13.4% Urea Nitrogen	
4.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	15%
Sulfur (S)	5.0%
5.0% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.1% Water Soluble Iron	
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
*7.3% Slowly Available Nitrogen from Methylene Ureas.	
Chlorine (Cl)	11.0%

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen	25%
4.3% Ammoniacal Nitrogen	
2.7% Water Insoluble Nitrogen*	
13.4% Urea Nitrogen	
4.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	15%
Sulfur (S)	5.0%
5.0% Combined Sulfur (S)	
Iron (Fe) Actual	1.0%
0.1% Water Soluble Iron	
Chlorine (Cl)	11.0%
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
*7.3% Slowly Available Nitrogen from Methylene Ureas.	

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

WARNING

Hazard Statements:

Causes eye irritation.
May be harmful if inhaled

Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling.
If in eyes, rinse with water and seek medical attention if symptoms persist.
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
If eye irritation persists: Get medical advice/attention.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen turf.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.6 lbs per 1,000 sq. ft. or 156.5 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.0 lbs per 1,000 sq. ft. or 87 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	0.9#N/1,000 sq. ft.	0.5#N/1,000 sq. ft.
Lebanon Turf	4½	4
Andersons AccuPro	L	H
Earthway Rotary	15	14
Gandy	27	24
Lely	5½ II	4 II
Lesco (letter/numeric)	i/19	G/15
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	4½	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.
To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 0.98 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.43 kg N/100 m ² .	0.22 kg N/100 m ² .
Lebanon Turf	4½	4
Andersons AccuPro	L	H
Earthway Rotary	15	14
Gandy	27	24
Lely	5½ II	4 II
Lesco (letter/numeric)	i/19	G/15
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	4½	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>



F699

22-37090

i/19



LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight