



ferti•lome®

St. Augustine Weed & Feed

15-0-4

**Use On St. Augustine, Zoysia, Centipede,
& Carpetgrass Lawns Only**

- Controls Crabgrass, Sedge, Spotted Spurge & Paspalum when applied before these weeds germinate
- Fertilize lawns & kill listed weeds in one easy step!
- Do not use on Bluegrass, Fescue, Bent, Bahia, Rye, Bermuda or Dichondra lawns

Dandelion
Pennywort
Chickweed
Sandbur
Henbit
Common Purslane

Burclover
Florida Betony
Florida Pusley
Creeping Charlie
Plantain
Lippia

Black Medic
Evening Primrose
Rabbit Tobacco
Annual Sowthistle
Blue Toadflax
Cranesbill

Dollarweed
Oxalis
Shepherdspurge
Virginia Pepperweed
Red Sorrel
Stinging Nettle

Wild Mustard
Dock
Rescuegrass
Pigweed
Venus Lookingglass

ACTIVE INGREDIENTS:

Atrazine: (2-chloro-4-ethylamino-6-isopropylamino-s-triazine) .. 0.79%
Related Compounds..... 0.02%

OTHER INGREDIENTS:..... 99.19%

TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

NET WEIGHT 12.8 LBS. (5.80 KG)

ferti•lome® St. Augustine Weed & Feed

ferti•lome® St. Augustine Weed & Feed

13915

ferti•lome®

**St. Augustine
Weed & Feed**



ferti•lome®

St. Augustine Weed & Feed

Use on St. Augustine, Zoysia, Centipede, and Carpetgrass Lawns Only.

| FIRST AID | |
|--------------------------------|--|
| If Swallowed: | <ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If On Skin Or Clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If In Eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue. • Call a poison control center or doctor for treatment advice. |
| If Inhaled: | <ul style="list-style-type: none"> • Move victim to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |

Have the product container or label with you when calling a poison control center or doctor, or going for help. You may also contact 1-800-222-1222 or 1-800-900-4044 for emergency medical treatment information.

NOTE TO PHYSICIAN: There is no specific antidote for Atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry or activated charcoal may be considered.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly after handling or using. Keep children and pets off treated areas until material is washed into the soil.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

NON-TARGET ORGANISM AND ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCT, OR FOR RESEARCH PURPOSES.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not enter or allow others to enter until dusts have settled. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. This product may be watered in immediately after application. Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry. Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments. Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).

1. WHAT IS THIS PACKAGE FOR?

This product is especially formulated to promote deep green, thick turf growth and eliminate unsightly weeds.

2. HOW DO YOU USE IT?

To prevent growth of Crabgrass, Sedge, Spotted Spurge, and Paspalum, apply in January, February, or March before these weeds germinate. These weeds will not be controlled if they are present in the lawn at the time of application. Apply in Fall to prevent Winter weeds such as Henbit, Chickweed, Rabbit Tobacco and Evening Primrose.

To kill other listed weeds, apply anytime weeds are present and actively growing. Repeat in 2 months if needed.

If soil is dry, water thoroughly the day before treatment. Treat entire lawn evenly. Immediately after application, water to wash particles off grass blades. **NOTE:**

Do not apply under the branch spread (root-zone) area of trees and shrubs in compact, slowly draining soils (particularly in the Dallas/Ft. Worth area). Do not use on Centipede or Carpetgrass lawns during the period of Winter dormancy (approximately October 15 to March 15), except in Florida. Do not make applications to St. Augustine, Zoysia, Centipede or Carpetgrass lawns containing other grasses. Do not apply to newly seeded Zoysia, Centipede or Carpetgrass lawns until after the new grass has been mowed at least twice. Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. Do not apply on bedding plants or ornamental trees and shrubs.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

Green, luxurious weed free lawns can be expected with this effective, easy-to-use product.

APPLICATION EQUIPMENT AND INSTRUCTIONS

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. Due to the wide swath, use broadcast lawn spreaders in open areas only; avoid trees, shrubs and flowers. To prevent over application and possible injury, spreader passes should be spaced an adequate distance apart to assure uniform coverage.

| SPREADER SETTINGS | |
|---|-----------------------------------|
| Spreader | Lbs. of product per 1,000 sq. ft. |
| | |
| Scotts®/Republic Accugreen (Drop) | 6 ¾ |
| Scotts®/Republic Speedy Green (Broadcast) | 6 ¼ |
| ferti-lome®/Earthway Ev-N-Spred (Broadcast) | 20 |

15-0-4 GUARANTEED ANALYSIS

| | |
|---|-------|
| Total Nitrogen (N) | 15% |
| 15% Urea Nitrogen* | |
| Soluble Potash (K ₂ O) | 4% |
| Boron (B) | 0.02% |
| Copper (Cu)..... | 0.05% |
| 0.05% Water Soluble Copper (Cu) | |
| Iron (Fe)..... | 1.0% |
| 0.1% Water Soluble Iron (Fe) | |
| Manganese (Mn) | 0.05% |
| 0.05% Water Soluble Manganese (Mn) | |
| Zinc (Zn)..... | 0.05% |
| 0.05% Water Soluble Zinc (Zn) | |

Derived From: Urea, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Iron Oxide, Manganese Sulfate, Zinc Sulfate.

* 4.5% Slowly Available Nitrogen derived from Polymer Coated Sulfur Coated Urea

F370

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep out or reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL: If empty: Non-refillable container. Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

Note: To the extent consistent with applicable law, Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

6F13915

13915

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 7401-336

EPA Est. No. 7401-TX-1

Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

ferti•lome®

St. Augustine Weed & Feed

