



DIMENSION™ 0.125% Plus Turf and Ornamental Fertilizer

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

A Selective Preemergent Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and other listed Weeds in Established Lawns and Ornamental Turfgrasses and Residential Landscapes.

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York this product may be applied only by commercial applicators at a minimum of 200 lb (0.25 lb of active ingredient) per acre (4.6 lb per 1,000 sq ft) and no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk Counties of New York is prohibited.

| | |
|-------------------------|--|
| Active Ingredient: | Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester.....0.125% |
| Other Ingredients | 99.875% |
| Total | 100.000% |

This product contains 0.0625 pound of the active ingredient dithiopyr per 50 pound bag.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

| FIRST AID | |
|--|---|
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 day or night, for emergency treatment information. | |

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

| ENVIRONMENTAL HAZARDS |
|---|
| This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions |

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

| Directions For Use |
|---|
| It is a violation of federal law to use this product in a manner inconsistent with its labeling. |
| STORAGE AND DISPOSAL |
| Do not contaminate water, foodstuffs, feed or seed by storage or disposal. |
| PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area. |
| PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically re-processed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. |
| CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. |

Use Directions for Turf

LESCO Dimension 0.125% Plus Turf and Ornamental Fertilizer provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury. Sweep any product that lands on driveway, sidewalk, or street back onto lawn and garden.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed.

| Cool-Season Grasses | |
|-----------------------|-----------------------------|
| Common Name | Scientific Name |
| bentgrass, creeping † | <i>Agrostis stolonifera</i> |
| bluegrass, Kentucky | <i>Poa pratensis</i> |
| fescue, fine †† | <i>Festuca rubra</i> |
| fescue, tall | <i>Festuca arundinacea</i> |
| ryegrass, perennial | <i>Lolium perenne</i> |

| Warm-Season Grasses | |
|---------------------|--------------------------------|
| Common Name | Scientific Name |
| bahiagrass | <i>Paspalum notatum</i> |
| bermudagrass | <i>Cynodon dactylon</i> |
| buffalograss ††† | <i>Buchloe dactyloides</i> |
| carpetgrass | <i>Axonopus affinis</i> |
| centipedegrass | <i>Eremochloa ophiuroides</i> |
| kikuyugrass | <i>Pennisetum clandestinum</i> |
| seashore paspalum | <i>Paspalum vaginatum</i> |
| St. Augustinegrass | <i>Stenotaphrum secundatum</i> |
| zoysiagrass | <i>Zoysia japonica</i> |

† Use of this product on certain varieties of Creeping Bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of Creeping Bentgrass have been tested. Do not apply this product to Colonial Bentgrass (*Agrostis tenuis*).

†† Use of this product on certain varieties of Fine Fescue may result in undesirable turf injury. The following Fine Fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of Fine Fescue have been tested.

††† Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: it is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

Precautions for Reseeding, Overseeding, or Sprigging

- Reseeding, overseeding or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation, and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (slit seeders) designed to provide good contact between soil and seed.

Precautions for Control of Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from areas treated with this product.

| |
|---|
| 18-0-3 |
| GUARANTEED ANALYSIS |
| TOTAL NITROGEN (N)18.00% |
| 12.00% Urea Nitrogen* |
| SOLUBLE POTASH (K ₂ O).....3.00% |
| DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash |
| CHLORINE (Cl) Max.....2.30% |
| *4.00% Slowly available Nitrogen from Polymer Sulfur-Coated Urea |

Application Directions for Turf

Apply LESCO Dimension® 0.125% Plus Turf and Ornamental Fertilizer with equipment that provides a uniform particle distribution. Calibrate equipment prior to usage. Avoid streaking, skips, or excess overlap during application.

Preemergence Application Rates, Frequency and Timing

For preemergence grass and broadleaf weed control, apply LESCO Dimension® 0.125% Plus Turf and Ornamental Fertilizer as single or sequential application at 200 to 400 pounds (0.25 to 0.5 lb active ingredient) per acre. Applicators may choose to make a single application or sequential applications at 200 to 400 pounds per acre at 5 to 10 week intervals based on one or more of the factors listed below. Avoid streaking, skips, or excess overlap during application.

- Length of residual weed control desired
- Height of turf (lower cut turf may require higher use rates)
- History and success of weed control at the application site (use higher listed application rates if herbicide treatment history is unknown or weed control was poor with previous applications)
- Exposure to high temperatures and heavy rainfall or irrigation (this will shorten the residual preemergence performance)
- On turf sites adjacent to hard surfaces including driveways, sidewalks, and parking lots where residual activity may be reduced
- Some target weed species (such as but not limited to *Poa annua*, goosegrass and sandbur) will require higher use rates

Tips for Improved Control

- For best results, apply this product within 2 to 3 days after mowing and delay mowing again for 2 to 3 days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation with 30 days after application.

Maximum Use Rates (Applies to all Turfgrass Uses)

- **Do not** apply more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per application.
- Do not apply more than 6 times per year **and do not** apply more than 1200 lb (1.5 lb active ingredient) per acre (27.6 lb per 1000 sq ft) per year if using split or sequential applications.
- To minimize the potential for turfgrass injury, do not make additional applications within five weeks of a previous application.
- In the state of New York, this product may be applied only by commercial applicators at no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Control of Crabgrass

Preemergence and Early Postemergence Applications from Late Winter through Summer

- This product provides "pre-emergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass at use rates from 200 to 400 pounds of product (0.25 to 0.5 lb ai) per acre.
- This product also provides "early post-emergence" control of crabgrass in late spring or early summer if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early post-emergence crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

Fall Applications

This product can also be applied in the late summer through early fall (late August through November) at use rates from 200 to 400 pounds per acre to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

Weeds Controlled or Suppressed

Spring Applications

When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds:

| | |
|------------------------|---------------------------------|
| Grasses | |
| barley | <i>Hordeum</i> spp. |
| barnyardgrass | <i>Echinochloa crus-galli</i> |
| bluegrass, annual | <i>Poa annua</i> |
| brome | <i>Bromus</i> spp. |
| crabgrass, large | <i>Digitaria sanguinalis</i> |
| crabgrass, smooth | <i>Digitaria ischaemum</i> |
| crabgrass, Southern | <i>Digitaria ciliaris</i> |
| crowfootgrass † | <i>Dactyloctenium aegyptium</i> |
| dallisgrass (seedling) | <i>Paspalum dilatatum</i> |
| goosegrass | <i>Eleusine indica</i> |
| foxtail, green | <i>Setaria verdi</i> |
| foxtail, yellow | <i>Setaria faberi</i> |
| kikuyugrass † | <i>Pennisetum clandestinum</i> |
| oats, wild | <i>Avena fatua</i> |
| ryegrass | <i>Lolium</i> spp. |
| (annual & perennial) | |
| sandbur | <i>Cenchrus</i> spp. |
| smutgrass | <i>Sporobolus indicus</i> |
| Broadleaf Weeds | |
| bittercress † | <i>Cardamine</i> spp. |
| carpetweed | <i>Mollugo verticillata</i> |
| chickweed † | <i>Stellaria</i> spp. |
| geranium, Carolina † | <i>Geranium carolinianum</i> |
| henbit | <i>Lamium</i> spp. |
| knottweed, prostrate | <i>Polygonum aviculare</i> |
| lespedeza, common † | <i>Lespedeza striata</i> |
| maestail | <i>Conyza canadensis</i> |
| medic, black | <i>Medicago lupulina</i> |
| mustard | <i>Brassica</i> spp. |

| | |
|----------------------|----------------------------------|
| oxalis, buttercup | <i>Oxalis pes-caprae</i> |
| pineappleweed † | <i>Matricaria matricarioides</i> |
| pigweed, redroot | <i>Amaranthus retroflexus</i> |
| parsley-piert † | <i>Alchemilla arvensis</i> |
| purslane, common | <i>Portulaca oleracea</i> |
| rocket, London | <i>Sisymbrium irio</i> |
| shepherdspurse | <i>Capsella bursa-pastoris</i> |
| speedwell, corn † | <i>Veronica arvensis</i> |
| spurge, garden | <i>Euphorbia hirta</i> |
| spurge, prostrate | <i>Euphorbia humistrata</i> |
| spurge, spotted | <i>Euphorbia maculata</i> |
| woodsorrel, creeping | <i>Oxalis corniculata</i> |
| woodsorrel, yellow | <i>Oxalis stricta</i> |

† **Suppression only**
Fall Applications

When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabgrass in the following spring.

| | |
|--------------------|----------------------------------|
| bluegrass, annual | <i>Poa annua</i> |
| bittercress | <i>Cardamine</i> spp. |
| chickweed | <i>Stellaria</i> spp. |
| henbit | <i>Lamium</i> spp. |
| geranium, Carolina | <i>Geranium carolinianum</i> |
| parsley-piert | <i>Alchemilla arvensis</i> |
| Pineappleweed | <i>Matricaria matricarioides</i> |
| Shepherdspurse | <i>Capsella bursa-pastoris</i> |

WEED RESISTANCE MANAGEMENT

LESCO® Dimension™ 0.125% Plus Turf and Ornamental Fertilizer, which contains the active ingredient dithiopyr is a Group 3 herbicide based on the mode of action classification system of the Weed Science Society of America. Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance. The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.
- Apply full rates of LESCO® Dimension™ 0.125% Plus Turf and Ornamental Fertilizer for the most difficult to control weed in the field at the specified time to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent.
- Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective modes of action for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 3 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-Group 3 herbicides.
- Avoid making more than two sequential applications of LESCO® Dimension™ 0.125% Plus Turf and Ornamental Fertilizer and any other Group 3 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields to reduce weed seed production.

Net Weight: 50 lb (22.7 kg) **SGN 240**
EPA REG. NO. 62719-483-10404
EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-002, 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1 EC.
 First letter of lot code indicates manufacturing site. **F1560**
 Distributed by: **LESCO, Inc.**
 1385 East 36th Street
 Cleveland, OH 44114-4114

702574 (p. 1)



DIMENSION™ 0.125% Plus Turf and Ornamental Fertilizer

APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granulates in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

| Rate | Calibration Settings | | |
|--------|------------------------------|------------------|--|
| | lbs. Product per 1,000 sq ft | lbs A.1 per acre | |
| Low | 4.6 | 0.25 | |
| Medium | 7 | 0.38 | |
| High | 9.2 | 0.5 | |

| Brand and Model of Spreader | Calibration Settings (lb/1000 sq ft)* | | |
|-----------------------------|---------------------------------------|--------------|------------|
| | Low (4.6) | Medium (7.0) | High (9.2) |
| LESCO | 16 | 19 | 23 |
| PermaGreen | 16 | 19 | 23 |
| Cyclone/Z-Spray | 4.25 | 4.75 | 5.25 |

[*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product]

Use Directions for Landscape Ornamentals

LESCO Dimension 0.125% Plus Turf and Ornamental Fertilizer provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. This product may be applied in ornamental gardens, parks, golf courses and residential areas where ornamental plants are grown for aesthetic purposes. Apply this product with a properly calibrated spreader that will provide uniform particle distribution. Follow instructions in the "Application Instructions and Equipment" section above.

Use Rates

This product may be applied as a single application or split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded.

| Pounds of Product/Acre | Pounds of Product/1000 sq ft | Pounds of Product/100 sq ft † |
|------------------------|------------------------------|-------------------------------|
| 400 | 9.2 | 0.92 (14.7 oz) |

†Rate based on an area of 100 sq ft for treatment of small areas.

Maximum Use Rates

- Do not apply more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per application.
- Do not apply more than 6 times per year **and do not** apply more than 1200 lb (1.5 lb active ingredient per acre (27.6 lb per 1000 sq ft) per year if using split or sequential applications.
- To minimize the potential for plant injury, do not make additional applications within five weeks of a previous application.
- **In the state of New York, this product may be applied only by commercial applicators at no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.**

Application Timing and Recommendations

LESCO Dimension 0.125% Plus Turf and Ornamental Fertilizer is effective as a preemergence herbicide, but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds, and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

Tips for Improved Weed Control

- The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

Use Precautions

- To avoid foliar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage.

Use Restrictions

- Do not apply this product directly to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes, or disease.
- Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.
- Apply this product only to established ornamentals.
- Do not use on food producing trees and ornamentals.

Weeds controlled or Suppressed

Used as directed, LESCO Dimension 0.125% Plus Turf and Ornamental Fertilizer will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination.

Refer to "Use Directions for Turf" section for a complete listing of weeds controlled or suppressed.

This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, only treat a few plants in a limited area and observe for tolerance under local growing conditions prior to large-scale use..

| Common Name | Botanical Name | Tolerant Cultivars |
|--------------------|--|--|
| abella, dwarf | <i>Abelia X grandiflora</i> | Nana |
| ajuga | <i>Ajuga reptans</i> | bronze bronze beauty |
| almond, flowering | <i>Prunus glandulosa</i> | |
| apple † | <i>Malus pumila</i> | |
| arborvitae | <i>Thuja occidentalis</i> | nigra pyramidalis smaragh techny woodwardii |
| arborvitae, golden | <i>Thuja orientalis</i> | |
| aster, Chinese | <i>Callistephus chinensis</i> | dwarf queen |
| ash, green | <i>Fraxinus pennsylvanica</i> | |
| ash, mountain | <i>Sorbus aucuparia</i> | |
| ash, purple | <i>Fraxinus americana</i> | |
| azalea | <i>Rhododendron spp.</i> | brilliant buccaneer carror chimes (belgian) elsie lee exbury fashion hardijzer beauty hershey red higasa hinocrimson holland (hybrid) marion lee northern lights orange cup orchid lights snow southern charm |
| azalea, flame | <i>Rhododendron calendulaceum</i> | |
| azalea, kirishima | | |
| bamboo, heavenly | | |
| barberry | <i>Berberis thunbergii</i> | aurea dwarf pigmy green kobold pygmy red rose glow |
| barberry, purple | | atropurpurea |
| basket flower | <i>Gaillardia grandiflora</i> | |
| bearberry (common) | <i>Arctostaphylos uva-ursi</i> | Massachusetts |
| bee balm | <i>Monarda didyma</i> | |
| begonia | <i>Begonia spp.</i> | |
| birch, river | <i>Betula nigra</i> | |
| blackeyed Susan | <i>Rudbeckia hirta</i> | goldstrum |
| blanket flower | <i>Gaillardia spp.</i> | |
| blueberry † | <i>Vaccinium spp.</i> | bluecrop blue jay jersey north blue northland |
| bottlebrush | <i>Callistemon citrinus</i> | |
| boxwood, Japanese | | japonica |
| boxwood, weller | <i>Buxus sempervirens</i> | |
| broom | <i>Cytisus spp.</i> <i>Genista pilosa</i> | moonlight Vancouver gold |
| bugle carpet | | |
| camellia | <i>Camellia japonica</i> <i>Camellia sasanqua</i> | debutante matholiana supreme chansonette |
| candy tuft | <i>Iberis spp.</i> | snow white |
| carex, variegated | <i>Carex</i> | |
| cedar, red | <i>Juniperus virginiana</i> | |
| celosia | <i>Celosia spp.</i> | |
| centaura | <i>Centaurea montana</i> | |
| cockscorn, plumosa | <i>Celosia cristata</i> | scarlet plumosa |
| coleus | <i>Coleus blumei</i> | red kewpie |
| columbine | <i>Aquilegia spp.</i> | |
| copper leaf | <i>Acalypha wilkesiana</i> | |
| coreopsis | <i>Coreopsis spp.</i> | moonbeam |
| corn flower | <i>Centaurea spp.</i> | |
| cotoneaster | <i>Cotoneaster apiculatus</i> | |
| coyotebrush | <i>Baccharis pilularis</i> | |
| cycads | <i>Cycads revoluta</i> | |
| cypress, bald | <i>Taxodium distichum</i> | |

| | | |
|--------------------------|--|--|
| cypress, Italian | <i>Cupressus sempervirens</i> | glauca |
| cypress, Japanese false | <i>Chamaecyparis obtusa</i> | gracilis |
| cypress, leyland | <i>Cupressocyparis leylandii</i> | |
| daffodil | <i>Narcissus spp.</i> | King Alfred |
| daylily | <i>Hemerocallis spp.</i> | aztec gold bright yellow (hybrid) single gold (evergreen) wilsonis yellow |
| dianthus (sweet william) | <i>Dianthus spp.</i> | |
| delphinium | <i>Delphinium spp.</i> | magic fountain |
| dogwood | <i>Cornus florida</i> | |
| dogwood, American | <i>Cornus sericea</i> | flavimarimaea |
| douglas fir | <i>Pseudotsuga menziesii</i> | |
| dusty miller | <i>Senecio cineraria</i> | |
| elm, drake | <i>Ulmus parvifolia</i> | |
| euonymus | <i>Euonymus fortunei</i> | argenteo-variegata auereo-marginata colorata emerald gaiety emerald en gold gold edge gold princess silver king tricolor vegetus |
| fan palm, European | <i>Chamaerops humilis</i> | |
| fan palm, Mexican | <i>Washingtonia robusta</i> | |
| fern (various) | <i>Asparagus spp.</i> | |
| fescue | <i>Festuca glauca</i> | |
| fetterbush | <i>Leucothoe fontanesiana</i> | rainbow |
| figus | <i>Ficus retusa</i> | nitidia |
| fir fraser | <i>Abies fraseri</i> | |
| forsythia | <i>Forsythia X intermedia</i> | Arnold dwarf bronzensis dwarf lynwood gold meadowlark weeping |
| | <i>Spring Glory</i> | |
| fuchsia | <i>Fuchsia spp.</i> | |
| galium | <i>Galium ordoatum</i> | |
| gardenia | <i>Gardenia jasminoides</i> | mystery radicans |
| geranium | <i>Pelargonium X hortorum</i> | |
| gum | <i>Eucalyptus citriodora</i> | |
| hawthorn | <i>Crataegus spp.</i> | cockspur white crimson cloud enchantress Jack Evans Washington white |
| heather, twisted | <i>Erica cinerea</i> | Mediterranean pink |
| hemlock, Canada | <i>Isuga canadensis</i> | |
| hibiscus | <i>Hibiscus spp.</i> | blue bird brilliant hula girl |
| holly | <i>Ilex spp.</i> | blue boy blue girl burfordii china girl compacta forsteri hellerie Japanese northern beauty needlepoint Nellie R. Stevens Savannah |
| | <i>Ilex X attenuata</i> | |
| holly, Chinese | <i>Ilex cornuta</i> | |
| holly, Japanese | <i>Ilex crenata</i> | |
| holly, yaupon | <i>Ilex vomitoria</i> | |
| honeysuckle | <i>Lonicera japonica</i> | claveyis dwarf halliana tatarian Canadian white zebelli red hosta |
| hosta | <i>Hosta spp.</i> <i>Hosta lancifolia</i> | albo marginata |
| ice plant | <i>Carpobrotus edulis</i> | |
| impatiens | <i>Impatiens spp.</i> | |
| iris | <i>Iris spp.</i> | dwarf blue wedgewood |
| ivy, English | <i>Hedera helix</i> | bulgaria thorndale |
| jasmine, Asian | <i>Trachelospermum asiaticum</i> | |
| juniper | <i>Juniperus spp.</i> | arcadia armstrong bar harbor |
| | <i>Juniperus horizontalis</i> | blue chip blue pacific blue rug |

| | | |
|---------------------|---------------------------------------|--|
| | <i>Juniperus horizontalis</i> | blue star broadmoor buffalo Calgary carpet emerald sea emerald spreader endora compacta fruitlandi green gold tip hetzi hughes Manhattan blue parsoni pfitzeriana plumosa prince of Wales procumbens dwarf San Jose sargent blue sargent green scandia scopolorum moon glow scopolorum skyrocket spartan |
| | <i>Juniperus chinensis</i> | tamariscifolia |
| | <i>Juniperus sabina</i> | weberi youngstown yukon belle |
| king palm | <i>Archontophoenix cunninghamiana</i> | |
| laurel, Australian | <i>Pittosporum tobira</i> | |
| laurel, mountain | <i>Kalmia latifolia</i> | |
| leucothoe | <i>Leucothoe fontanesiana</i> | |
| ligustrum, Japanese | <i>Ligustrum japonicum</i> | |
| lily, African | <i>Agapanthus africanus</i> | albus Peter Pan |
| lily, African blue | | |
| lily of the valley | <i>Piensi japonica</i> | Mt. Fire |
| lilyturf | <i>Liriope muscari</i> | evergreen giant lilac beauty majestic monroe white variegata |
| liriope, green | <i>Liriope spicata</i> | |
| maple, Japanese | <i>Acer japonicum</i> | |
| maple, Norway | <i>Acer platanoides</i> | |
| maple, red † | <i>Acer rubrum</i> | |
| maple, silver | <i>Acer saccharinum</i> | |
| maple sugar † | <i>Acer saccharum</i> | |
| marigold | <i>Tagetes patula</i> | honeycomb variegata wheeleris dwarf |
| mock orange † | <i>Philadelphus spp</i> | golden snowflake double white |
| mondo grass | <i>Ophiopogon japonicus</i> | |
| moss rose | <i>Portulaca grandiflora</i> | sunnyside |
| myrtle, crape | <i>Lagerstroemia indica</i> | faurei langer muskogee standard pink |
| myrtle, wax | <i>Myrica cerifera</i> | |
| nandina | <i>Nandina domestica</i> | compacta nana |
| narcissus | <i>Narcissus spp.</i> | |

Net Weight: 50 lb (22.7 kg)

SGN 240

EPA REG. NO. 62719-483-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-002, 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1 EC.

First letter of lot code indicates manufacturing site.

F1560

Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

702574-PG2



DIMENSION™ 0.125% Plus Turf and Ornamental Fertilizer

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, only treat a few plants in a limited area and observe for tolerance under local growing conditions prior to large scale use.

| Common Name | Botanical Name | Tolerant Cultivars |
|-------------------------|---|--|
| oak, laurel | <i>Quercus laurifolia</i> | |
| oak, pin | <i>Quercus palustris</i> | |
| oak, red | <i>Quercus rubra</i> | |
| oak, southern | <i>Quercus virginiana</i> | |
| oak, willow | <i>Quercus phellos</i> | |
| oleander | <i>Nerium oleander</i> | hardy red petite pink Sister Agnes |
| osteospermum | <i>Osteospermum fruticosum</i> | wirrligig |
| pachysandra | <i>Pachysandra terminalis</i> | |
| palm, bangalow | | |
| pampas grass | <i>Cortaderia selloana</i> | |
| pansy | <i>Viola x wittrockiana</i> | |
| paper flower | <i>Bougainvillea glabra</i> | Barbara karst |
| peach † | <i>Prunus persica</i> | |
| periwinkle, dwarf | <i>Vinca minor</i> | |
| petunia | <i>Petunia X hybrida</i> | picoti |
| photinia, red tip | <i>Photinia X fraseri</i> | |
| plenis | <i>Pieris japonica</i> | |
| pine, Australian | <i>Pinus nigra</i> | |
| pine, Japanese black | <i>Pinus thunbergiana</i> | |
| pine, loblolly | <i>Pinus taeda</i> | |
| pine, longleaf | <i>Pinus palustris</i> | |
| pine, mugo | <i>Pinus mugo</i> | |
| pine, Scotch | <i>Pinus sylvestris</i> | |
| pine, slash | <i>Pinus elliotii</i> | |
| pine, Swiss mt. | <i>Pinus mugo</i> | |
| pine, Virginia | <i>Pinus virginiana</i> | |
| pine, white | <i>Pinus strobus</i> | |
| pineapple, guava † | <i>Feijoa sellowiana</i> | |
| plenis | <i>Pieris taiwanensis</i> | |
| pittosporum, Japan | | |
| potentilla | <i>Potentilla nepalensis</i> <i>Potentilla fruticosa</i> | abbotswood |
| privet | <i>Ligustrum japonicum</i> | golden vicary regal texanum wax yellow tipped |
| privet, glossy | <i>Ligustrum lucidum</i> | |
| pyracantha | <i>Pyracantha koidzumii</i> | gnome lalandei victory |
| queen palm | <i>Arecastrum rammanzoffianum</i> | |
| quince, Japanese † | | |
| rhododendron | <i>Rhododendron spp</i> | album Cunningham white fashion hardy pjm purple gem silvery pink |
| rhododendron, carolina | <i>Rhododendron carolinianum</i> | |
| rhododendron, catawba | <i>Rhododendron catawbiense</i> | |
| rockcress | <i>Arabis spp.</i> | snowcap |
| rhodie max (rosebay) | <i>Rhododendron maximum</i> | |
| rose † | <i>Rosa banksiae</i> | luta |
| rosemary † | <i>Rosmarinus officinalis</i> | |
| rosemary, bog | <i>Andromeda polifolia</i> | nana |
| salvia | <i>Salvia faninacea</i> | rhea |
| sedum | <i>Sedum spurium</i> | dragon blood red red carpet yellow |
| snapdragon | <i>Antirrhinum spp.</i> | |
| sourwood | <i>Oxydendrum arboreum</i> | |
| spiraea | <i>Astilbe X arendsii</i> | fanall |
| spiraea | <i>Spiraea spp.</i> | Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white |
| spiraea, garland | <i>Spiraea spp.</i> | |
| spruce, black hills | <i>Picea glauca</i> | |
| spruce, Colorado blue | <i>Picea pungens</i> | |
| spruce, norway | <i>Picea abies</i> | |
| spruce, white | <i>Picea glauca</i> | conica |
| sweetflag, grassy-le | | |
| sweetgum | <i>Liquidambar styraciflua</i> | |
| sycamore | <i>Platanus occidentalis</i> | |
| tree fern (tiki fern) | <i>Asparagus virgatus</i> | |
| trumpet flower, evening | <i>Gelsemium sempervirens</i> | |

| | | |
|--------------------|--|--|
| tulip | <i>Tulip, spp</i> | apeldoorn |
| verbena, shrub | <i>Lantana sellowiana</i> | |
| vervain | <i>Verbena spp.</i> | St. Paul |
| viburnum | <i>Viburnum spp.</i> | American cranberry bush arrowood common snowball |
| vinca (periwinkle) | <i>Vinca minor</i> | |
| windmill palm | <i>Trachycarpus fortunei</i> | |
| xylosma | <i>Xylosma congestum</i> | |
| yarrow | <i>Achillea spp.</i> | |
| yaupon | <i>Ilex vomitoria</i> | dwarf |
| yew | <i>Taxus cuspidata</i> <i>Taxus X media</i> | denisiformis |

† Ornamental species only: Do not use on food producing trees and ornamentals.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.htm>

LESCO is a registered trademark and the sweeping design is a trademark of LESCO, Inc. DIMENSION is a registered trademark of Dow AgroSciences. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. N:\Regulatory\WPI\Regul Private\MstrLb\ERIEVIEW ADDRESS\Landscape Style Master Lbs (070208) Rev 12/23/19 KT

Net Weight: 50 lb (22.7 kg)

SGN 240

EPA REG. NO. 62719-483-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-002, 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1 EC.

First letter of lot code indicates manufacturing site.

F1560

Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

#702574