



MERIT® 0.2% Plus Turf Fertilizer

For fertilization, insect control in turfgrass, landscape ornamentals and sod farms.
Fertilizer with MERIT® Insecticide

50 lbs. treats 14,200 sq. ft. (0.33 acre) at the 150 lbs. product/acre - LOW RATE (3.5 lbs/1,000 sq. ft.)
50 lbs. treats 11,100 sq. ft. (0.25 acre) at the 200 lbs. product/acre - HIGH RATE (4.5 lbs/1,000 sq. ft.)

ACTIVE INGREDIENT:
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine0.2%

OTHER INGREDIENTS:99.8%

Total100.0%

STOP – Read the label before use KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

For MEDICAL and TRANSPORTATION Emergencies ONLY

Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-321-5325.

Have a product container or label with you when calling a poison control center or doctor or going for treatment.

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
No specific antidote is available. Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WPS USES: Applicators and Other Handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR part 170) – in general, agricultural plant uses e.g., use in sod farms, must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Non-WPS USES: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR part 170) – in general, only agricultural plant uses are covered by the WPS, must wear:

- Shirt and pants
- Gloves
- Shoes plus socks

USER SAFETY RECOMMENDATION:

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

10-0-0

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)10.00%

10.00% Urea Nitrogen

DERIVED FROM: Urea.

*

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-refillable container. Do not re-use or refill this container. Offer for recycling if available. Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration or if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed.

Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above.

Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

APPLICATION TO TURFGRASS

The use of this product will provide plant nutrients as well as providing control of selected insect pests. LESCO® MERIT® 0.2% Plus Turf Fertilizer can be used for the control of soil inhabiting pests of turfgrass, for example, Northern & Southern masked chafers, Cyclocephala borealis, C. immaculata, and/or C. lurida; Asiatic garden beetle, Maladera castanea; European chafer, Rhizotrogus majalis; Green June beetles, Cotinus nitida; May or June beetles, Phyllophaga spp.; Japanese beetle, Popillia japonica; Oriental beetle, Anomala orientalis; Billbugs, Sphenophorus spp.; Annual bluegrass weevil, Listronotus maculicollis. Black turfgrass ateniens, Ateniens spretulus; Aphodius spp.; crane flies, Tipula spp., and mole crickets, Scapteriscus spp.

The active ingredient in LESCO® MERIT® 0.2% Plus Turf Fertilizer can be used as directed on turfgrass, for example, in home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms.

The active ingredient in LESCO® MERIT® 0.2% Plus Turf Fertilizer has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically into the soil profile.

APPLICATION TO ORNAMENTALS

LESCO® MERIT® 0.2% Plus Turf Fertilizer can be used as a broadcast application on ornamentals in commercial and residential landscapes, as well as on landscaped areas of golf courses and other recreational areas. This is a systemic product, with the active ingredient being absorbed by the roots and translocated upward within the plant. Optimum control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation, or rainfall to place the active ingredient into the soil. Make application prior to anticipated pest infestation to achieve optimum levels of control.

Applications of LESCO® MERIT® 0.2% Plus Turf Fertilizer cannot exceed a total of 200 lb (0.4 lb of active ingredient) per acre per year.

RESTRICTION: Not for use in commercial greenhouses, or nurseries.

APPLICATION RATES			
CROP	PEST		DOSAGE
			LESCO® MERIT® 0.2% Plus Turf Fertilizer
Turf Grasses	Annual bluegrass weevil Billbugs Crane flies Larvae of white grubs, for example: Aphodius sp. Asiatic garden beetle Black turfgrass ateniens European chafer	Green June beetle Japanese beetle Northern masked chafer Oriental beetle Phyllophaga spp Southern masked chafer Sugarcane beetle	150 to 200 lb per acre or 3.5 to 4.5 lb per 1000 sq ft
	For optimum control, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.		
	Mole crickets Chinch bugs (suppression)		200 lb per acre or 4.5 lb per 1000 sq ft
For suppression of chinch bugs, make application prior to the hatching of the first instar nymphs. For control of mole crickets make application within one week of the peak egg hatch period. If application is not made prior to egg hatch, and adults or large nymphs are present and actively tunneling, apply LESCO® MERIT® 0.2% Plus Turf Fertilizer in combination with a remedial insecticide to minimize damage to turf.			
Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.			
USE RESTRICTIONS			
<ul style="list-style-type: none"> • Do not apply more than 200 lb (0.4 lb of active ingredient) per acre per year. • Do not apply to soils which are waterlogged or saturated, which will not allow penetration into the root zone of the plant. • For optimum control, irrigate if rainfall does not occur within 24 hours after application to move the active ingredient through the thatch. • On golf courses, irrigate treated areas following application. • For optimum control, mow turf or lawn area only after irrigation or rainfall has occurred so that uniformity of application will not be affected. • Avoid runoff or puddling of irrigation water following application. 			

APPLICATION RATES FOR USE ON ORNAMENTALS

SITE	PEST	DOSAGE
		LESCO® MERIT® 0.2% Plus Turf Fertilizer
Landscape Plantings Including: Trees, Shrubs, Evergreens, Flowers, Foliage, Plants, Ground Covers	White grub larvae for example : Japanese beetle larvae, Chafers, Phyllophaga spp., Asiatic garden- beetle, Oriental beetle	3.5 to 4.5 lb per 1,000 sq ft
Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.		
USE RESTRICTIONS		
<ul style="list-style-type: none"> • Do not apply this product, by any application method, to linden, basswood, or other Tilia species. • Do not apply more than 200 lb (0.4 lb of active ingredient) per acre per year. • For optimum control, irrigate if rainfall does not occur, or incorporate mechanically, within 24 hours after application to move the active ingredient into the soil. • Avoid runoff or puddling of irrigation water following application. • Avoid application of LESCO® MERIT® 0.2% Plus Turf Fertilizer to areas which are water-logged or saturated, which will not allow penetration into the root zone of the plant. • Do not apply to vegetable gardens. • Refer to label sections on application equipment and spreader settings. 		

APPLICATION EQUIPMENT

Apply LESCO® MERIT® 0.2% Plus Turf Fertilizer uniformly over the area being treated with typical granular application equipment. Both drop-type and rotary-type spreaders may be used to apply LESCO® MERIT® 0.2% Plus Turf Fertilizer. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

SPREADER SETTING FOR LESCO MERIT 0.2 PLUS TURF FERTILIZER

The following chart can be used as a guide in establishing an accurate rate of LESCO® MERIT® 0.2% Plus Turf Fertilizer. The suggested settings provide an application rate of 3.5 lb per 1000 sq ft, or 150 lb per acre. Application rates will vary dependent upon speed, equipment wear and condition of spreader. Carefully calibrate all equipment used prior to commercial use of this product.

Spreader	SETTINGS
LESCO® – All Rotary Models	#14
PermaGreen	14
Cyclone®/ Spyker® /Z-Spray	4
Vicon (LESCO® Pendulum)	22
Lely®	4 II

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of LESCO, INC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of LESCO, INC. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LESCO, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

An extended Spreader Setting listing can be found at: www.lesco.com.

MERIT is a registered trademark of Bayer.
LESCO and the sweeping design is a trademark of LESCO Inc.
Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.
Vicon is a registered trademark of Ransomes-Cushman-Ryan.
PermaGreen is a registered trademark of PermaGreen Supreme, Inc.
Lely is a registered trademark of C. Van Der Lely N.V.

(022509) Rev. 8/4/22 VT

Net Weight: 50 lb (22.7 kg) SGN 240

EPA REG. NO. 432-1349-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-004 (BAN), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1-EC, 92626-TN-1, 49991-TX-2, 93243-CT-1, 93243-TX-1.

First letter of lot code indicates manufacturing site.

F1560
Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

#293008