



ECHELON™ 0.30% Plus Fertilizer ::OPTI

PRODIAMINE	GROUP	3	HERBICIDE
SULFENTRAZONE	GROUP	14	HERBICIDE

For Preemergence and early postemergence Control of Crabgrass, selected annual grasses, sedges and annual broadleaf weeds in Turf Sites - Residential and Institutional Lawns, Athletic Fields, Commercial Sod Farms, Golf Course Fairways and Roughs, Roadsides, Utility right of ways, Railways and Industrial areas.

Contains LESCO® Poly Plus® OPTI™ Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENTS:

Sulfentrazone:.....	0.1%
Prodiamine:.....	0.2%
OTHER INGREDIENTS:	99.7%
TOTAL:	100.0%

This product contains 0.003 pounds of active ingredient (0.001 lb ai sulfentrazone and 0.002 lb ai prodiamine) per pound of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See other panels for additional precautionary information.

FIRST AID

If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

MEDICAL HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

For Technical Support or information regarding the use of this product, call 1-800-347-4272.

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

- Users should remove clothing/PPE immediately if product gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants and aquatic organism in adjacent areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface water advisory: Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas over-lying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product may only be used in accordance with the Application Rates listed on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

WEED RESISTANCE MANAGEMENT

For resistance management, please note that this product contains both a Group 3 and a Group 14 herbicide. Any weed population may contain plants naturally resistant to Group 3 and/or Group 14 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 3 and/or Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout area before herbicide application for weed identification and growth state.
- Scout after herbicide application to monitor weed populations for early signs of resistance development.

Indicators of possible herbicide resistance include:

- 1) Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds
 - 2) A spreading patch of non-controlled plants of a particular weed species
 - 3) Surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your local extension specialist or university researchers for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.

11-0-5	
GUARANTEED ANALYSIS	
TOTAL NITROGEN (N)	11.00%
11.00% Urea Nitrogen	
SOLUBLE POTASH (K ₂ O).....	5.00%
IRON (Fe).....	3.00%
0.03% Water Soluble Iron (Fe)	
MANGANESE (Mn)	1.00%
0.01% Water Soluble Manganese (Mn)	
DERIVED FROM: Polymer Coated Urea, Urea, Muriate of Potash, Iron Sulfate, Manganese Sulfate.	
CHLORINE (Cl) Max.	3.75%
*3.30% Slowly available Urea Nitrogen from Polymer Coated Urea.	
<p>FLORIDA APPLICATIONS: For use on golf courses or other athletic turf in FL, we recommend you follow SL191, "Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Based on Mehlich I Extractant", available on the internet at http://edis.ifas.ufl.edu/SS404.</p> <p>FLORIDA APPLICATIONS: For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at: http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf</p>	

PRODUCT INFORMATION

LESCO Echelon™ 0.30% Plus Fertilizer is a granular herbicide on fertilizer carrier formulation containing 0.003 pounds of active ingredient per pound of formulated product. The mode of action of this product involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings. This product is a selective pre-emergence and post-emergence herbicide that may only be used to control crabgrass, annual grass and broadleaf weeds and sedges in turf sites including residential and institutional lawns, athletic fields, commercial sod farms, golf course fairways and roughs, roadsides, utility right of ways, railways and industrial areas.

SPECIFIC USE INSTRUCTIONS IN TURFGRASS

ESTABLISHED TURF

This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences. Broadcast this product with suitable application equipment to ensure uniform coverage over the treatment area. A drop or rotary spreader designed to apply granular herbicide or insecticide is the preferred application of this product. Calibrate application equipment prior to application. Consult manufacturer operation manual for proper calibration and operation of application equipment.

LESCO Echelon™ 0.30% Plus Fertilizer application may cause temporary discoloration to exposed leaf surfaces on certain cultivars of zoysiagrass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply this product on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.

LESCO Echelon™ 0.30% Plus Fertilizer has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties and/or cultivars have been evaluated. Turfgrass managers desiring to treat newly re-leased varieties should first apply this product to a small area prior to treatment of larger areas.

APPLICATION RATES

Turfgrass Use Rate Conversion				
lbs product per 1000 sq ft	lbs Sulfentrazone per acre	lbs Prodiamine per acre	lbs total ai per acre	lbs product per acre
1.91	0.08	0.17	0.25	83
2.87	0.125	0.25	0.375	125
4.4	0.19	0.38	0.57	190
5.7	0.25	0.50	0.75	250
8.6	0.375	0.750	1.125	375

When used as directed under the conditions described, apply this product at 83 - 375 lbs product per acre (1.91 - 8.6 lbs product per 1,000 sq ft) per application. The maximum single and annual application rate is 375 lbs product per acre (equivalent to 0.375 lbs sulfentrazone and 0.750 lbs prodiamine per acre) per calendar year. Refer to Table 1 for the application rates for each grass type.

Table 1. Application Rates for Tolerant grasses.

GRASS TYPE	USE RATES	
	Do not exceed maximum use rates per turf species in a single application	
Cool Season Grasses	lbs product/1000 sq ft	lbs product/acre
Bentgrass, creeping ¹ , Fescue, fine ² , (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>)	1.91 - 2.87 ³	83 - 125 ³
Bluegrass, Kentucky (<i>Poa pratensis</i>), Fescue, tall ² (<i>Festuca arundinacea</i>)	4.4 - 5.7 ³	190 - 250 ³
Sequential Application for Mixed Cool Season Turf/Cool Season Blends consisting of:		
Fescue, fine ² , (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>) Bluegrass, Kentucky (<i>Poa pratensis</i>), Fescue, tall ² (<i>Festuca arundinacea</i>)	Initial application 2.87	Initial application 125
	Follow-up Application 1.91 - 2.87 30 to 60 days later	Follow-up Application 83 to 125 lbs 30 to 60 days later

Warm Season Grasses		
Bahiagrass (<i>Paspalum notatum</i>) Buffalograss (<i>Buchloe dactyloides</i>) Carpentergrass (<i>Axonopus affinis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kikuyugrass (<i>Pennisetum clandestinum</i>) Seashore Paspalum (<i>Paspalum vaginatum</i>) Zoysiagrass (<i>Zoysia japonica</i>)	4.4 - 5.7 ³	190 - 250 lbs
Bermudagrass (<i>Cynodon dactylon</i>) & hybrids	5.7 - 8.6 ³	250 - 275 lbs
<ol style="list-style-type: none"> 1. If bentgrass is mowed lower than 0.5 inches do not apply this product. 2. Use of this product on certain varieties of Cheering Fine Fescue or Tall Fescue cultivars may result in undesirable injury 3. This product may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate per turf type or 375 lbs per acre (equivalent to 0.375 lbs sulfentrazone and 0.750 lbs prodiamine per acre), whichever is lower. 		

Table 1A. Application for St. Augustinegrass
Application of this product to St. Augustinegrass may cause temporary discoloration to exposed leaf surfaces of St. Augustinegrass. Treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. Do not apply this product to St. Augustinegrass if temporary discoloration cannot be accepted.

Warm Season Grass	lbs product/ 1000 sq ft	lbs product/acre
St. Augustinegrass	Initial Application 4.4	Initial Application 190
	Follow-up Application 4.4	Follow-up Application 190
	45 to 60 days later	45 to 60 days later

Preemergence Control of Annual, Biennial & Perennial Broadleaf Weeds and sedges
This product, applied alone or in recommended tank mixtures, will provide control of the following weeds for pre-emergence applications.

Net Weight: 50 lb (22.7 kg) SGN 240

EPA REG. NO. 279-3322-10404

Rev. 3/3/22 VT

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-004 (BAN), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1-EC, 92626-TN-1, 49991-TX-2, 93243-CT-1

First letter of lot code indicates manufacturing site.
F1560
Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

#702690-PG1



ECHELON™ 0.30% Plus Fertilizer

PRODIAMINE	GROUP	3	HERBICIDE
SULFENTRAZONE	GROUP	14	HERBICIDE



Table 2. Weed Controlled with Preemergence Application

Common Name	Scientific Name
Barnyardgrass	(<i>Echinochloa crus-galli</i>)
Bluegrass annual ¹	(<i>Poa annua</i>)
Carpetweed	(<i>Mollugo verticillata</i>)
Carolina Geranium	<i>Geranium carolinianum</i>
Chickweed, common ²	(<i>Stellaria media</i>)
Chickweed, mouseear (from seed)	(<i>Cerastium vulgatum</i>)
Crabgrass (large, Smooth) ²	(<i>Digitaria spp.</i>)
Crowfootgrass	(<i>Dactyloctenium aegyptium</i>)
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)
Foxtails, Annual	(<i>Setaria spp.</i>)
Goosegrass ¹	(<i>Elymus indica</i>)
Hairy Bittercress	<i>Cardamine hirsute</i>
Henbit ²	(<i>Lamium amplexicaule</i>)
Itchgrass	(<i>Rottboellia exaltata</i>)
Knotweed, prostrate	(<i>Polygonum aviculare</i>)
Kochia	(<i>Kochia scoparia</i>)
Kyllinga, green	(<i>Kyllinga brevifolia</i>)
Kyllinga, false	(<i>Kyllinga gracillima</i>)
Lambsquarters, common	(<i>Chenopodium album</i>)
Lovegrass	(<i>Eragrostis spp.</i>)
Nutsedge, yellow	(<i>Cyperus esculentus</i>)
Panicum (Texas, Fall)	(<i>Panicum spp.</i>)
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)
Purslane, Common	(<i>Portulaca oleracea</i>)
Pusley, Florida	(<i>Richardia scabra L.</i>)
Sedge, globe	(<i>Cyperus globulosus</i>)
Sedge, cylindrical	(<i>Cyperus retrosus</i>)
Sedge, Sunnam	(<i>Cyperus surinamensis</i>)
Sedge, Texas	(<i>Cyperus polystachyos</i>)
Shephardspurse ²	(<i>Capsella bursa-pastoris</i>)
Signalgrass, broadleaf	(<i>Brachiaria platyphylla</i>)
Speedwell	(<i>Veronica spp.</i>)
Sprangletop	(<i>Leptochloa spp.</i>)
Spurge, prostrate	(<i>Euphorbia humistrata</i>)
Witchgrass	(<i>Panicum capillare L.</i>)
Woodsorrel, yellow (from Seed)	(<i>Oxalis stricta</i>)

1. Poa annua is a winter annual. Applications of this product should be made in August or September to establish, non overseeded turf before Poa Annua germinates. Consult local Cooperative Extension Service for more specific application timings for your local area since August and September are approximate timings.
2. Apply this product in late summer, fall or winter before weed germinates for control.
3. Crabgrass control in cool season grasses from fall application: Areas where the ground is subject to winter freezes, this product may be applied in the fall at rates listed in Table 1 (83-250 lbs per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.
4. In areas of extended growing season and heavy pressure weed control will be enhanced by second application of this product 60-90 days after initial application.

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds and sedges

This product, applied alone or in recommended tank mixture, will provide control of the following weeds for post emergence applications.

Table 3. Weeds Controlled or Suppressed When Application Made Postemergence to Weeds

Common Name	Scientific Name	Controlled	Suppressed
Bittercress	(<i>Cardamine spp.</i>)		X
Black medic	(<i>Medicago lupulina</i>)		X
Buttercups	(<i>Ranunculus spp.</i>)		X
Carolina Geranium	<i>Geranium carolinianum</i>	1	
Carpetweed	(<i>Mollugo verticillata</i>)		X
Chickweed, common	(<i>Stellaria media</i>)	1	
Chickweed, mouseear	(<i>Cerastium vulgatum</i>)		X
Cinquefoil	(<i>Potentilla spp.</i>)		X
Clover	(<i>Trifolium spp.</i>)		X
Crabgrass (Large and Smooth)	(<i>Digitaria spp.</i>)	2	
Cudweed	(<i>Gnaphalium spp.</i>)		X
Dandelion	(<i>Taraxacum officinale</i>)		X
Dock, Curly	(<i>Rumex crispus</i>)		X
Evening primrose	(<i>Oenothera biennis</i>)		X
Fiddleneck	(<i>Amsinckia spp.</i>)		X
Filaree	(<i>Erodium spp.</i>)		X
Goldenrod	(<i>Solidago spp.</i>)	1	
Goosegrass	(<i>Elymus indica</i>)	2	
Ground ivy	(<i>Glechoma hederacea</i>)	1	
Henbit	(<i>Lamium amplexicaule</i>)		X
Knotweed, prostrate	(<i>Polygonum aviculare</i>)		X
Kochia	(<i>Kochia scoparia</i>)	1	
Lambsquarters, common	(<i>Chenopodium album</i>)		X
Lespedeza, common	(<i>Lespedeza striata</i>)		X
Mallow, common	(<i>Malva neglecta</i>)		X
Nutsedge, yellow	(<i>Cyperus esculentus</i>)	1	
Parsley piert	(<i>Alchemilla arvensis</i>)	1	
Pigweed Redroot	(<i>Amaranthus retroflexus</i>)	1	
Pigweed, Tumble	(<i>Amaranthus albus</i>)	1	
Pineapple weed	(<i>Matricaria matricarioides</i>)		X
Plantain, buckhorn	(<i>Plantago lanceolata</i>)	1	
Puncture weed	(<i>Tribulus terrestris</i>)		X
Purslane, Common	(<i>Portulaca oleracea</i>)		X
Pusley, Florida	(<i>Richardia scabra L.</i>)	1	
Redweed	(<i>Melochia corchorifolia</i>)		X

Common Name	Scientific Name	Controlled	Suppressed
Rocket, London	(<i>Sisymbrium lino</i>)		X
Smartweed, Pennsylvania	(<i>Polygonum pennsylvanicum</i>)	1	
Sorrel, Red	(<i>Rumex acetosella</i>)		X
Speedwell	(<i>Veronica spp.</i>)	1	
Spurge, (annuals)	(<i>Euphorbia spp.</i>)	1	
Spurge, prostrate	(<i>Euphorbia humistrata</i>)	1	
Spurge, spotted	(<i>Euphorbia maculata</i>)	1	
Star of Bethlehem	(<i>Ornithogalum umbellatum</i>)	1	
Velvet leaf	(<i>Abutilon theophrasti</i>)		X
Violet, wild	(<i>Viola pratensis</i>)		X
Wild garlic	(<i>Allium vineale</i>)		X
Wild onion	(<i>Allium canadense</i>)		X
Woodsorrel, creeping	(<i>Oxalis corniculata</i>)		X
Woodsorrel, yellow	(<i>Oxalis stricta</i>)	1	

1. Weeds are suppressed at lower label rates (<250 lb product/acre). For optimum control apply rates of at least 250 lb product per acre in a single application. Do not exceed the application rate specified for the turf species in Table 1.
2. This product controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.

APPLICATION TO RESEEDED, OVER- SEEDED OR SPRIGGED AREAS

Reseeding, overseeding or sprigging of treated areas within three to twelve (3-12) months after application of this product could inhibit the establishment of desirable grasses

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

Table 4. Overseeding Intervals after Product Applications

Total Amount of Product Applied lb product/ Acre	Interval (Months Before Overseeding)		
	North	Transition	South
83 – 125	3	3	3
190 – 250	4	4	4
375	6	5	5

SOD PRODUCTION

Sod should be established for at least six (6) months before an application of this product.

Do not apply this product within three (3) months of harvest.

TURFGRASS USE RESTRICTIONS

- Do not apply to golf course putting greens or tees.
- Do not use on turf grasses other than those listed on this label.
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply directly to landscape ornamentals or ornamental beds.
- Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.
- Do not apply to creeping bentgrass if mowing height is less than 0.5 inch.
- Do not harvest treated sod before 90 days after application.
- Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.
- Do not apply this product through any type of irrigation system.
- Do not apply by air.

ADDITIONAL USE INSTRUCTIONS

Railroad Rights-of-Way

Controls many weeds and maintains bare ground on railroad rights-of-way, including railroad yards, railroad crossings and railroad bridge abutments.

Highway, Roadside, Pipeline and Utility Rights-of-Way

Controls many weeds and maintains bare ground in highway, roadside, pipeline and utility rights-of-way. Such areas would include guardrails, road shoulders, electric utility substations, pipeline pumping stations, around electric transmission towers, around distribution line poles plus other areas where complete vegetation control is desired.

Industrial Areas, Fence Rows and Other Non-crop Sites

Controls weeds and maintains bare ground in industrial areas including production facilities, tank farms, storage areas, parking areas, lumber yards, airports, military installations, along fence rows, and in similar non-crop sites where complete vegetation control is needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Reseal open bag by folding top down and securing. Store in a cool, dry place and avoid excess heat.

Spill: In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. **Confine spills. Call CHEMTREC (Transportation and spills): (800) 424-9300.** **To confine spill:** If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Paper and Plastic Bags: Non-refillable. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, if allowed by state or local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfc.org/metals.htm>

Poly Plus® OPTI™ is comprised of Polymer Coated Urea. LESCO, Poly Plus and Echelon are registered trademarks and the sweeping design is a trademark of LESCO, Inc. PermaGreen is a registered trademark of PermaGreen Supreme, Inc. OPTI™ is a registered trademark of Turf Care Supply Corp. Vicon is a registered trademark of Ransomes-Cushman-Ryan. Rev. 3/3/22 VT

Net Weight: 50 lb (22.7 kg) SGN 240

EPA REG. NO. 279-3322-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-004 (BAN), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1-EC, 92626-TN-1, 49991-TX-2, 93243-CT-1

First letter of lot code indicates manufacturing site.

F1560
Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

#702690-PG2