# The Andersons Turf Products Fertilizer with 0.2% MERIT<sup>®</sup> Insecticide **21-0-4**

- · For fertilization and insect control in turfgrass and landscape ornamentals and sod farms.
- Fertilizer with MERIT<sup>®</sup> Insecticide.
- Contains MERIT<sup>®</sup> Insecticide.

**Guaranteed Analysis** 

Total Nitrogen (N)	21.00%
Soluble Potash (K <sub>2</sub> O).	4.00%
Plant nutrients derived from urea, polymer coated sulfur coated urea and potassium chloride. Chlorine (CI) Max	3.40%
*10.50% Slowly available urea nitrogen from polymer coated sulfur coated urea.	F002091

ACTIVE INGREDIENT:	
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	0.2%
OTHER INGREDIENTS:	99.8%
TOTAL:	100.0%

FPA Reg No 432-1349-9198 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Underlined letter is first letter used in run code on bag

MERIT is a registered trademark of Baver

### STOP - READ THE LABEL BEFORE USE. **KEEP OUT OF REACH OF CHILDREN** CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le hava sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

#### For <u>MEDICAL</u> and <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-800-331-2867

FIRST AID		
If swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>	
If on skin or clothing:	Take off contaminated clothing.     Rinse skin immediately with plenty of water for 15-20 minutes.     Call a poison control center or doctor for treatment advice.	
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	

Note to Physician: No specific antidote is available. Treat the patient symptomatically

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

#### Personal Protective Equipment (PPE):

WPS USES: Applicators and Other Handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR part 170) - in general, agricultural plant uses e.g., use in sod farms, must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPF separately from other laundry

Non-WPS USES: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR part 170) - in general, only agricultural plant uses are covered by the WPS, must wear: Shirt and pants

· Gloves

· Shoes plus socks

#### Aphodius sp This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas Asiatic garde Black turfgras where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. European ch DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For optimum Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area Mole crickets during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation. Chinch bugs (suppression) AGRICULTURAL USE REQUIREMENTS Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. USE RESTRICTIONS Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is Coveralls Chemical-resistant gloves made of any waterproof material Shoes plus socks NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep SITE children and pets off treated area until dry. Landscape Plantings APPLICATION TO TUREGRASS The use of this product will provide plant nutrients as well as providing control of selected insect pests. This product can be used for the control of soil Including Trees, Shrubs, inhabiting pests of turfgrass; for example northern and southern masked chafers, Cyclocephala borealis, C. immaculata, and/or C. lurida; Asiatic garden beetle, Maladera castanea; European chafer, Ahizotrogus majalis; Green June beetle, Colinus nitida; May or June beetles, Phyllophaga spp.; Japanese beetle, Popillia japonica; Oriental beetle, Anomala orientalis; Euetheola humulis rugiceps; sugarcane beetle, Euetheola humulis rugiceps, billbugs, Sphenophorus spp.; annual bluegrass weevil, Listronotus maculicollis; Black turfgrass ataenius, Ataenius spretulus; Aphodius spp.; crane flies Tipula Evergreens, Flowers, Foliage Plants. Ground Covers spp., and mole crickets, *Scapteriscus* spp. Use this product as directed on turgrass, including home lawns, school grounds, nursing homes and hospital grounds, city, county, state, municipal and federal grounds, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, churches, cemeteries, parks, playgrounds, athletic fields and sod farms. USE RESTRICTIONS The active ingredient in this product has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch. · Do not apply to vegetable gardens. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically into the soil profile. APPLICATION TO ORNAMENTALS This product can be used as a broadcast application on ornamentals in commercial and residential landscapes, as well as on landscaped areas of golf courses and other recreational areas. This is a systemic product, with the active ingredient being absorbed by the roots and translocated upward within the plant. Optimum control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation or rainfall to place the active ingredient into the soil. Make application prior to anticipated pest infestation to achieve optimum levels of control. Applications of this product cannot exceed a total of 200 lb. (0.4 lb. of active ingredient) per acre per year RESTRICTION: Not for use in commercial greenhouses, or nurseries. APPLICATION EQUIPMENT Apply this product uniformly over the area being treated with typical granular application equipment. Both drop-type and rotary-type spreaders may be used to apply this product. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately. SPREADER SETTINGS FOR THIS PRODUCT The following chart can be used as a guide in establishing an accurate rate of this product. The suggested settings provide an application rate of 3.5 lb per 1,000 sq.ft. or 150 lb per acre. Application rates will vary dependent upon speed, equipment wear and condition of spreader. Carefully or buver calibrate all equipment used prior to use of this product. 50 lbs. treats 14,200 sg. ft. (0.33 acre) at the 150 lbs. product/acre - LOW RATE (3.5 lbs/1,000 sg. ft.)

USER SAFETY RECOMMENDATION

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

below the mean high water mark. Do not contaminate water when disposing of equipment washwaters

· Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

Users should:

and change into clean clothing.

ENVIRONMENTAL HAZARDS

			SPREADER SETTINGS	
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE 0.3 lb. ai/A	HIGH RATE 0.4 lb. ai/A
Andersons Model 2000 / AccuPro 2000 (cone 4) Andersons 2000 SR / SR-2000 (cone 4)	3 mph	10 feet	L	М
The Andersons LCO-1000	3 mph	10 feet	L	М
Scotts R-8A (cone 4)	3 mph	10 feet	K 1/4	L 1/4
Lesco 020093 (pattern 0.25)	3 mph	8 feet	E 1/2	F 1/2
Lesco 705698 (pattern 0.25)	3 mph	8 feet	13	15
Lely Models - WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern III-A	4.5 mph	32 feet	5 1/2	6
Vicon (03 Series)	4.5 mph	28 feet	24	29

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

IMPORTANT: READ BEFORE USE Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of The Andersons, Inc.. All such risks shall be assumed by the use

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE ANDERSONS, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of The Andersons, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE ANDERSONS, INC. DISCLAIMS ANY liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ANDERSONS, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT

APPLICATION RATES			
PES	т	DOSAGE THE ANDERSONS TURF PRODUCTS FERTILIZER WITH 0.2% MERIT INSECTICIDE 21-0-4	
Annual bluegrass weevil Billbugs Crane flies Larvae of white grubs, for example: <i>Aphodius</i> sp. Asiatic garden beetle Black turfgrass ataenius European chafer	Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer Sugarcane beetle	150 to 200 lbs. per acre or 3.5 to 4.5 lbs. per 1,000 sq. ft.	
For optimum control, make application	prior to egg hatch of the target pest.	Be sure to read "APPLICATION EQUIPMENT" Section of this label.	
		200 lbs per acre	

CROP

Billbugs Crane flies

Turf Grasses

For suppression of chinch bugs, make application prior to the hatching of the first instar nymphs. For control of mole crickets make application within one week of the peak egg hatch period. If application is not made prior to egg hatch, and adults or large nymphs are present and actively tunneling, apply this product in combination with a remedial insecticide to minimize damage to turf.

4.5 lbs. per 1,000 sq. ft.

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application

Do not apply more than 200 lb (0.4 lb of active ingredient) per acre per year.

Do not apply to soils which are waterlogged or saturated, which will not allow penetration into the root zone of the plant.
 For optimum control, irrigate if rainfall does not occur within 24 hours after application to move the active ingredient through the thatch.
 On golf courses, irrigate treated areas following application.

For optimum control, mow turf or lawn area only after irrigation or rainfall has occurred so that uniformity of application will not be affected.
 Avoid runoff or puddling of irrigation water following application.

APPLICATION RATES FOR USE ON ORNAMENTALS		
	PEST	DOSAGE THE ANDERSONS TURF PRODUCTS FERTILIZER WITH 0.2% MERIT INSECTICIDE 21-0-4
	White grub larvae	3.5 to 4.5 lbs. per 1,000 sq. ft.
	For example: Japanese beetle larvae, chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle	

Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.

Do not apply this product, by any application method, to linden, basswood, or other Tilia species.
Do not apply more than 200 lb (0.4 lb of active ingredient) per acre per year.

For optimum control, irrigate if rainfall does not occur, or incorporate mechanically, within 24 hours after application to move the active ingredient into the soil.
 Avoid runoff or puddling of irrigation water following application.

· Avoid application of this product to areas which are water-logged or saturated, which will not allow penetration into the root zone of the plant.

Refer to label sections on application equipment and spreader settings.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Handling: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration or if allowed by the state and local authorities, by burning. If burned, stay out of smoke,

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

Manufactured by: The Andersons, Inc P.O. Box 119 Maumee, OH 43537

## Net Wt. 50 Lbs. (22.68 kg)



**APT211MR5.40**